

Signed combinatorial interpretations in algebraic combinatorics

Colleen Robichaux
UCLA

joint work with Igor Pak

FPSAC 2025, Hokkaido University
July 21, 2025

Combinatorial Structure Constants

Consider a basis $B = \{\xi_\alpha\}$ in a ring R :

$$\xi_\alpha \xi_\beta = \sum_{\gamma \in A} c_{\alpha\beta}^\gamma \xi_\gamma$$

Here A is some combinatorial set (partitions, compositions, etc).

Example

Take $R = \mathbb{Z}[x_1, \dots, x_n]^{S_n}$ and $B = \{s_\lambda : \ell(\lambda) \leq n\}$:

$$s_\lambda s_\mu = \sum_{\nu} c_{\lambda\mu}^\nu s_\nu.$$

Here $c_{\lambda\mu}^\nu \in \mathbb{Z}_{\geq 0}$ are the *Littlewood–Richardson coefficients*.

A Central Motivation

Problem A

Determine a *non-cancellative* combinatorial rule to compute structure coefficients $c_{\alpha\beta}^{\gamma}$.

Why should such a rule exist?

A Central Motivation

Problem A

Determine a *non-cancellative* combinatorial rule to compute structure coefficients $c_{\alpha\beta}^{\gamma}$.

Why should such a rule exist?

Problem B

Determine a *cancellative* combinatorial rule to compute structure coefficients $c_{\alpha\beta}^{\gamma}$.

What is “combinatorial”?

A Formalization

Informal Definition

Counting X is in $\#P$ if there exists a set S with $|S| = |X|$, where checking $s \in S$ is poly-time.

Example: LR-coefficients $c_{\lambda,\mu}^{\nu}$

The LR-rule via ballot semistandard Young tableaux gives

$$c_{\lambda,\mu}^{\nu} = \#\text{ballot SSYT}(\nu/\lambda, \mu) \in \#P$$

A Formalization

Define **GapP** $:= \#P - \#P$.

Example: symmetric group characters χ_μ^λ

The Murnaghan–Nakayama rule via border strip tableaux shows

$$\chi_\mu^\lambda = \sum_{T \in \text{BST}(\lambda, \mu)} (-1)^{ht(T)} \in \text{GapP}$$

Problem B, reformulated

Determine a GapP rule for $c_{\alpha\beta}^\gamma$.

Central Takeaway

Combinatorial bases with integral structure constants have signed combinatorial formulas.

Central Takeaway

Combinatorial bases with integral structure constants have signed combinatorial formulas.

(**Problem B** is easily resolvable in most cases)

Key Ingredient: Möbius Inversion

Let $\mathcal{P} := (A, \prec)$ be a poset.

Hall's Theorem

Take $\eta : A^2 \rightarrow \mathbb{Z}$ unitriangular. Then the inverse $\rho : A^2 \rightarrow \mathbb{Z}$ of η has the form:

$$\rho(x, y) = \sum_{\ell=0}^h \sum_{\substack{\text{chains} \\ x \rightarrow z_1 \rightarrow \dots \rightarrow z_{\ell-1} \rightarrow y}} (-1)^\ell \eta(x, z_1) \cdot \eta(z_1, z_2) \cdot \dots \cdot \eta(z_{\ell-1}, y).$$

Proposition

Suppose $\mathcal{P}$ has polynomial height and the incidence function in $\mathcal{P}$ is poly-time computable. Then η is in GapP implies ρ is in GapP.


Schubert polynomials

Schubert polynomials give a $\mathbb{Z}$ -basis of $R = \mathbb{Z}[x_1, x_2, \dots, x_n]$:

$$\mathfrak{S}_w(x_1, \dots, x_n) = \sum_{P \in PD(w)} \mathbf{x}^{\text{wt}(P)} = \sum_{\alpha \in \mathbb{Z}_{\geq 0}^n} K_{c(w), \alpha} \mathbf{x}^\alpha.$$

Here $c(w) \in \mathbb{Z}_{\geq 0}^n$ is the **code** of $w \in S_n$.

Example: $\mathfrak{S}_w$ for $w = 21435$ via Bergeron–Billey '96



$$\text{so } \mathfrak{S}_w(x_1, x_2, x_3, x_4, x_5) = x_1^2 + x_1x_2 + x_1x_3.$$

Setup for $\mathfrak{S}_w$

Take $\mathcal{P} = (V_{n,k}, \trianglelefteq)$ to be the poset on $\alpha \in \mathbb{Z}_{\geq 0}^n$ with $|\alpha| = k$. Here $\trianglelefteq$ is dominance order. Then:

- $\mathcal{P}$ has polynomial height
- $K_{c(w),\alpha} \in \#P$
- $\mathfrak{S}_w(x_1, \dots, x_n) = \sum_{c(w) \trianglelefteq \alpha} K_{c(w),\alpha} \mathbf{x}^\alpha$
- diagonal terms $K_{c(w),c(w)} = 1$

Consider the inverse Schubert Kostka coefficients

$$\mathbf{x}^\alpha = \sum_{c(w) \trianglelefteq \alpha} K_{c(w),\alpha}^{-1} \mathfrak{S}_w(x_1, \dots, x_n).$$

Corollary

$$K_{c(w),\alpha}^{-1} \in \text{GapP}$$

Argument for $\mathfrak{S}_w$

$$\begin{aligned}\mathfrak{S}_u \cdot \mathfrak{S}_v &= \left(\sum_{c(u) \trianglelefteq \alpha} K_{c(u), \alpha} \mathbf{x}^\alpha \right) \cdot \left(\sum_{c(v) \trianglelefteq \beta} K_{c(v), \beta} \mathbf{x}^\beta \right) \\&= \sum_{c(u) \trianglelefteq \alpha} \sum_{c(v) \trianglelefteq \beta} \left(K_{c(u), \alpha} K_{c(v), \beta} \right) \mathbf{x}^{\alpha+\beta} \\&= \sum_{c(u) \trianglelefteq \alpha} \sum_{c(v) \trianglelefteq \beta} \sum_{\alpha+\beta \trianglelefteq c(w)} \left(K_{c(u), \alpha} K_{c(v), \beta} K_{c(w), \alpha+\beta}^{-1} \right) \mathfrak{S}_w \\&= \sum_w c_{uv}^w \mathfrak{S}_w.\end{aligned}$$

Corollary

Schubert structure coefficients $c_{uv}^w \in \text{GapP}$

Main Theorem

Theorem [Pak–R. '24]

Structure constants for the following families are in GapP:

- Symmetric Bases

- $s_\lambda, m_\lambda, e_\lambda, h_\lambda, p_\lambda$

- Quasisymmetric Bases

- $M_\alpha, F_\alpha, \mathfrak{S}_\alpha^*, S_\alpha$

- Polynomial Bases

- $\mathfrak{M}_\alpha, \mathfrak{F}_\alpha, \text{atom}_\alpha, \kappa_\alpha, \mathfrak{S}_w, \mathcal{L}_\alpha, \mathfrak{G}_w$

Using the same techniques, plethysm coefficients $a_{\lambda\mu}^\nu$ for $f_\lambda[g_\mu] = \sum_\nu a_{\lambda\mu}^\nu s_\nu$ are also in GapP for symmetric bases f_λ, g_μ .

Main Theorem – Extended

Theorem [Pak–R. '24]

Structure constants for the following are in GapP (or GapP/FP):

- Symmetric Bases

- $s_\lambda, m_\lambda, e_\lambda, h_\lambda, p_\lambda$
- $P_\lambda(x; \alpha), P_\lambda(x; t), P_\lambda(x; q, t)$ (GapP/FP)

- Quasisymmetric Bases

- $M_\alpha, F_\alpha, \mathfrak{S}_\alpha^*, \mathcal{S}_\alpha$
- $\Psi_\alpha, \Phi_\alpha, \mathfrak{p}_\alpha$ (GapP/FP)

- Polynomial Bases

- $\mathfrak{M}_\alpha, \mathfrak{F}_\alpha, \text{atom}_\alpha, \kappa_\alpha, \mathfrak{S}_w, \mathcal{L}_\alpha, \mathfrak{G}_w$


Here GapP/FP are functions f/g where $f \in \text{GapP}$ and g is poly-time computable (i.e. in FP).

What next? Improved signed formulas!

We construct $O(n^9)$ puzzle pieces $\mathcal{T}_n$. For $u, v, w \in S_n$, we build a parallelogram region $\Gamma(u, v, w)$. Using Knutson's recurrence '03:

Theorem (Pak–R. '25)

The number of signed puzzles of $\Gamma(u, v, w)$ using $\mathcal{T}_n$ is $c_{u,v}^w$.



Application

Using this signed puzzle rule, we find:

Corollary (Pak–R. '25)

Fix k , and let

$$\gamma_k(n) := \sum_{u,v,w \in S_n: \text{inv}(w)=k} c_{u,v}^w.$$

Then γ_k is a polynomial in n .

This follows from inequalities imposed by our rule + Erhart theory.

Conclusion

- Integral structure constants are in GapP
- Plethysm coefficients are in GapP
- Develop improved GapP formulas for your favorite combinatorial basis
- Use GapP formulas to illuminate existing structure, develop asymptotics, and approach $\#P$ formulas

Thank you!