

(P, ϕ) -TAMARI LATTICES

ADRIEN SEGOVIA (LACIM-UQAM)

Under the supervision of Samuele Giraudo and Hugh Thomas

FPSAC 2025

July 22, Hokkaido University

LACIM

Laboratoire d'algèbre, de combinatoire et
d'informatique mathématique



SUMMARY:

BACKGROUND



I) POSETS AND LATTICES

NEW
CONSTRUCTION



II) (P, ϕ) -TAMARI


III) HIGHER TORSION CLASSES:



ALGEBRAIC MOTIVATION

TYPE A AUSLANDER AND NAKAYAMA ALGEBRAS

I) POSETS AND LATTICES




GIAN-CARLO ROTA, ROBERT DILWORTH, GARRETT BIRKHOFF, ERNST SCHRÖDER

- A **LATTICE** $(L, \leq)$ IS A POSET SUCH THAT ALL $\{x, y\} \subseteq L$ HAVE A LEAST UPPER BOUND AND A GREATEST LOWER BOUND.
 MEET $x \wedge y$ JOIN $x \vee y$


THE **CHAINS** ARE THE TOTALLY ORDERED SUBSETS. THE **LENGTH** OF L IS $\ell(L) :=$ SIZE OF A MAXIMUM-LENGTH CHAIN.

BIGGER

 SMALLER



NOT A LATTICE



LATTICE


- A **LATTICE** $(L, \leq)$ IS A POSET SUCH THAT ALL $\{x, y\} \subseteq L$ HAVE A LEAST UPPER BOUND AND A GREATEST LOWER BOUND.
 $\swarrow$ MEET $x \wedge y$ $\searrow$ JOIN $x \vee y$

THE **CHAINS** ARE THE TOTALLY ORDERED SUBSETS. THE **LENGTH** OF L IS $\ell(L) :=$ SIZE OF A MAXIMUM-LENGTH CHAIN.


DENOTE $J\mathcal{I}(L)$ THE SET OF **JOIN-IRREDUCIBLES**: THOSE THAT COVER A UNIQUE ELEMENT. WE DEFINE $M\mathcal{I}(L)$ DUALY.

AN ORDER **IDEAL** (**FILTER**) IS THE SUBSET OF ELEMENTS BELOW (ABOVE) SOME ELEMENTS. $\mathcal{I}_p(x) := \{y \mid y \leq x\}$.


BIGGER
 $\updownarrow$
 SMALLER



NOT A LATTICE



LATTICE




- A **LATTICE** $(L, \leq)$ IS A POSET SUCH THAT ALL $\{x, y\} \subseteq L$ HAVE A LEAST UPPER BOUND AND A GREATEST LOWER BOUND. MEET $x \wedge y$ JOIN $x \vee y$

THE **CHAINS** ARE THE TOTALLY ORDERED SUBSETS. THE **LENGTH** OF L IS $\ell(L) :=$ SIZE OF A MAXIMUM-LENGTH CHAIN.

DENOTE $JI(L)$ THE SET OF **JOIN-IRREDUCIBLES**: THOSE THAT COVER A UNIQUE ELEMENT. WE DEFINE $MI(L)$ DUALY. AN ORDER **IDEAL** (**FILTER**) IS THE SUBSET OF ELEMENTS BELOW (ABOVE) SOME ELEMENTS. $I_p(x) := \{y \mid y \leq x\}$.

- L IS **JOIN-SEMIDISTRIBUTIVE** IF $\forall x, y, z \in L, x \vee y = x \vee z \Rightarrow x \vee (y \wedge z) = x \vee y$.

MEET-SEMIDISTRIBUTIVE IS DEFINED DUALY, **SEMIDISTRIBUTIVE** IF BOTH JOIN AND MEET-SEMIDISTRIBUTIVE.




- A **LATTICE** $(L, \leq)$ IS A POSET SUCH THAT ALL $\{x, y\} \subseteq L$ HAVE A LEAST UPPER BOUND AND A GREATEST LOWER BOUND. MEET $x \wedge y$ JOIN $x \vee y$

THE **CHAINS** ARE THE TOTALLY ORDERED SUBSETS. THE **LENGTH** OF L IS $\ell(L) :=$ SIZE OF A MAXIMUM-LENGTH CHAIN.


DENOTE $JI(L)$ THE SET OF **JOIN-IRREDUCIBLES**: THOSE THAT COVER A UNIQUE ELEMENT. WE DEFINE $MI(L)$ DUALY. AN ORDER **IDEAL** (**FILTER**) IS THE SUBSET OF ELEMENTS BELOW (ABOVE) SOME ELEMENTS. $I_p(x) := \{y \mid y \leq x\}$.

- L IS **JOIN-SEMIDISTRIBUTIVE** IF $\forall x, y, z \in L, x \vee y = x \vee z \Rightarrow x \vee (y \wedge z) = x \vee y$.
MEET-SEMIDISTRIBUTIVE IS DEFINED DUALY, **SEMIDISTRIBUTIVE** IF BOTH JOIN AND MEET-SEMIDISTRIBUTIVE.
- L IS **EXTREMAL** IF $\ell(L) = |JI(L)| = |MI(L)|$. JOIN-EXTREMAL


BIGGER
↑
↓
SMALLER




NOT A LATTICE



LATTICE



JOIN-SEMIDISTRIBUTIVE



SEMIDISTRIBUTIVE

- A **LATTICE** $(L, \leq)$ IS A POSET SUCH THAT ALL $\{x, y\} \subseteq L$ HAVE A LEAST UPPER BOUND AND A GREATEST LOWER BOUND. MEET $x \wedge y$ JOIN $x \vee y$





THE **CHAINS** ARE THE TOTALLY ORDERED SUBSETS. THE **LENGTH** OF L IS $\ell(L) :=$ SIZE OF A MAXIMUM-LENGTH CHAIN.

DENOTE $\text{JI}(L)$ THE SET OF **JOIN-IRREDUCIBLES**: THOSE THAT COVER A UNIQUE ELEMENT. WE DEFINE $\text{MI}(L)$ DUALY. AN ORDER **IDEAL** (**FILTER**) IS THE SUBSET OF ELEMENTS BELOW (ABOVE) SOME ELEMENTS. $\mathcal{I}_p(x) := \{y \mid y \leq x\}$.


- L IS **JOIN-SEMIDISTRIBUTIVE** IF $\forall x, y, z \in L, x \vee y = x \vee z \Rightarrow x \vee (y \wedge z) = x \vee y$.

MEET-SEMIDISTRIBUTIVE IS DEFINED DUALY, **SEMIDISTRIBUTIVE** IF BOTH JOIN AND MEET-SEMIDISTRIBUTIVE.

- L IS **EXTREMAL** IF $\ell(L) = |\text{JI}(L)| = |\text{MI}(L)|$. **JOIN-EXTREMAL**
- $a \in L$ IS **LEFT MODULAR** IF $\forall b < c, (b \vee a) \wedge c = b \vee (a \wedge c)$. IF L HAS A LEFT MODULAR MAXIMAL CHAIN, THEN L IS LEFT MODULAR.




- THE **ORDER COMPLEX** $\Delta(P)$ IS THE SIMPLICIAL COMPLEX WHOSE FACES ARE THE CHAINS OF P .
- IT IS **SHELLABLE** IF WE CAN ORDER THE FACETS $F_1, F_2, \dots, F_m$ SUCH THAT FOR ALL k , $\left(\bigcup_{1 \leq i < k} \langle F_i \rangle\right) \cap \langle F_k \rangle$ IS A UNION OF PURE COMPLEXES OF DIMENSION $\dim(\langle F_k \rangle) - 1$.



- THE **ORDER COMPLEX** $\Delta(P)$ IS THE SIMPLICIAL COMPLEX WHOSE FACES ARE THE CHAINS OF P .
- IT IS **SHELLABLE** IF WE CAN ORDER THE FACETS $F_1, F_2, \dots, F_m$ SUCH THAT FOR ALL k , $\left(\bigcup_{1 \leq i < k} \langle F_i \rangle\right) \cap \langle F_k \rangle$ IS A UNION OF PURE COMPLEXES OF DIMENSION $\dim(\langle F_k \rangle) - 1$.
- LET $\phi: \hat{O} = x_0 < x_1 < \dots < x_k = \hat{1}$ BE A CHAIN OF L . FOR $(j, m) \in \text{JI}(L) \times \text{MI}(L)$, $\begin{cases} \delta(j) := \min \{ i \mid j \leq x_i \} \\ \beta(m) := \max \{ i \mid m \geq x_{i-1} \} \end{cases}$.
 $\gamma_1(b < c) := \min \{ \delta(j) \mid j \in \text{JI}(L), j \leq c, j \not\leq b \}$ $\gamma'_1(b < c) := \max \{ i \mid c \wedge x_{i-1} \leq b \}$
 $\gamma_2(b < c) := \max \{ \beta(m) \mid m \in \text{MI}(L), m \geq b, m \not\geq c \}$ $\gamma'_2(b < c) := \min \{ i \mid b \vee x_i \geq c \}$

PROP [S.]: $(\gamma_1 = \gamma'_1 = \gamma_2 = \gamma'_2) \Leftrightarrow (\forall i, x_i \text{ IS LEFT MODULAR}).$


[Liu(99)]



- THE **ORDER COMPLEX** $\Delta(P)$ IS THE SIMPLICIAL COMPLEX WHOSE FACES ARE THE CHAINS OF P .
- IT IS **SHELLABLE** IF WE CAN ORDER THE FACETS $F_1, F_2, \dots, F_m$ SUCH THAT FOR ALL k , $\left(\bigcup_{1 \leq i < k} \langle F_i \rangle\right) \cap \langle F_k \rangle$ IS A UNION OF PURE COMPLEXES OF DIMENSION $\dim(\langle F_k \rangle) - 1$.
- LET $\phi: \hat{O} = x_0 < x_1 < \dots < x_k = \hat{1}$ BE A CHAIN OF L . FOR $(j, m) \in \text{JI}(L) \times \text{MI}(L)$, $\begin{cases} \delta(j) := \min \{i \mid j \leq x_i\} \\ \beta(m) := \max \{i \mid m \geq x_{i-1}\} \end{cases}$.
 $\gamma_1(b < c) := \min \{ \delta(j) \mid j \in \text{JI}(L), j \leq c, j \not\leq b \}$ $\gamma'_1(b < c) := \max \{ i \mid c \wedge x_{i-1} \leq b \}$
 $\gamma_2(b < c) := \max \{ \beta(m) \mid m \in \text{MI}(L), m \geq b, m \not\geq c \}$ $\gamma'_2(b < c) := \min \{ i \mid b \vee x_i \geq c \}$

PROP [S.]: $(\gamma_1 = \gamma'_1 = \gamma_2 = \gamma'_2) \Leftrightarrow (\forall i, x_i \text{ IS LEFT MODULAR}).$
[LIU(99')]

COROLLARY: [T.W. (199')] SEMIDISTRIBUTIVE EXTREMAL LATTICE ARE LEFT MODULAR.



● THE **ORDER COMPLEX** $\Delta(P)$ IS THE SIMPLICIAL COMPLEX WHOSE FACES ARE THE CHAINS OF P .

● IT IS **SHELLABLE** IF WE CAN ORDER THE FACETS $F_1, F_2, \dots, F_m$ SUCH THAT FOR ALL k , $\left(\bigcup_{1 \leq i < k} \langle F_i \rangle\right) \cap \langle F_k \rangle$ IS A UNION OF PURE COMPLEXES OF DIMENSION $\dim(\langle F_k \rangle) - 1$.

● LET $\phi: \hat{O} = x_0 < x_1 < \dots < x_k = \hat{1}$ BE A CHAIN OF L . FOR $(j, m) \in \text{JI}(L) \times \text{MI}(L)$, $\begin{cases} \delta(j) := \min \{i \mid j \leq x_i\} \\ \beta(m) := \max \{i \mid m \geq x_{i-1}\} \end{cases}$.


$$\gamma_1(b < c) := \min \{ \delta(j) \mid j \in \text{JI}(L), j \leq c, j \not\leq b \} \quad \gamma'_1(b < c) := \max \{ i \mid c \wedge x_{i-1} \leq b \}$$

$$\gamma_2(b < c) := \max \{ \beta(m) \mid m \in \text{MI}(L), m \geq b, m \not\geq c \} \quad \gamma'_2(b < c) := \min \{ i \mid b \vee x_i \geq c \}$$

PROP [S.]: $(\gamma_1 = \gamma'_1 = \gamma_2 = \gamma'_2) \Leftrightarrow (\forall i, x_i \text{ IS LEFT MODULAR}).$

[LIU(99)]

COROLLARY: [T.W. (199')] SEMIDISTRIBUTIVE EXTREMAL LATTICE ARE LEFT MODULAR.



● A **(JOIN-) CONGRUENCE UNIFORM** LATTICE IS ONE OBTAINED BY THE DOUBLING CONSTRUCTION FROM THE ONE ELEMENT LATTICE BY DOUBLING ONLY (LOWER PSEUDO-) INTERVALS.

● A **CONGRUENCE** ON L IS AN EQUIVALENCE RELATION α SUCH THAT $\left. \begin{matrix} x_1 \equiv x_2 [\alpha] \\ y_1 \equiv y_2 [\alpha] \end{matrix} \right\} \Rightarrow \begin{cases} x_1 \wedge y_1 \equiv x_2 \wedge y_2 [\alpha] \\ x_1 \vee y_1 \equiv x_2 \vee y_2 [\alpha] \end{cases}$.

● THE **ORDER COMPLEX** $\Delta(P)$ IS THE SIMPLICIAL COMPLEX WHOSE FACES ARE THE CHAINS OF P .

● IT IS **SHELLABLE** IF WE CAN ORDER THE FACETS $F_1, F_2, \dots, F_m$ SUCH THAT FOR ALL k , $\left(\bigcup_{1 \leq i < k} \langle F_i \rangle\right) \cap \langle F_k \rangle$ IS A UNION OF PURE COMPLEXES OF DIMENSION $\dim(\langle F_k \rangle) - 1$.

● LET $\phi: \hat{O} = x_0 < x_1 < \dots < x_k = \hat{1}$ BE A CHAIN OF L . FOR $(j, m) \in \text{JI}(L) \times \text{MI}(L)$, $\begin{cases} \delta(j) := \min \{i \mid j \leq x_i\} \\ \beta(m) := \max \{i \mid m \geq x_{i-1}\} \end{cases}$.


$$\gamma_1(b < c) := \min \{ \delta(j) \mid j \in \text{JI}(L), j \leq c, j \not\leq b \} \quad \gamma'_1(b < c) := \max \{ i \mid c \wedge x_{i-1} \leq b \}$$

$$\gamma_2(b < c) := \max \{ \beta(m) \mid m \in \text{MI}(L), m \geq b, m \not\geq c \} \quad \gamma'_2(b < c) := \min \{ i \mid b \vee x_i \geq c \}$$

PROP [S.]: $(\gamma_1 = \gamma'_1 = \gamma_2 = \gamma'_2) \Leftrightarrow (\forall i, x_i \text{ IS LEFT MODULAR}).$

[LIU(99)]

COROLLARY: [T.W. (199')] SEMIDISTRIBUTIVE EXTREMAL LATTICE ARE LEFT MODULAR.




● A **(JOIN-) CONGRUENCE UNIFORM** LATTICE IS ONE OBTAINED BY THE DOUBLING CONSTRUCTION FROM THE ONE ELEMENT LATTICE BY DOUBLING ONLY (LOWER PSEUDO-) INTERVALS.

● A **CONGRUENCE** ON L IS AN EQUIVALENCE RELATION α SUCH THAT $\left. \begin{matrix} x_1 \equiv x_2 [\alpha] \\ y_1 \equiv y_2 [\alpha] \end{matrix} \right\} \Rightarrow \begin{cases} x_1 \wedge y_1 \equiv x_2 \wedge y_2 [\alpha] \\ x_1 \vee y_1 \equiv x_2 \vee y_2 [\alpha] \end{cases}$.


THM [S.]: A CONGRUENCE UNIFORM LATTICE IS SHELLABLE IFF IT IS EXTREMAL.

II) (P, ϕ) -TAMARI




LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$,
 CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$.
 $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING
 FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.

LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$,
 CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$.
 $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING
 FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.




P


LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$, CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$. $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.




LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$, CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$. $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.




LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$, CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$. $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.






LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$, CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$. $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.




LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$, CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$. $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.



LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{p, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$, CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{p, \phi}$. $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.





LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$, CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$. $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.




ORDER FILTERS:

$\emptyset = 000$
 $\{(a,0)\} = 100$
 $\{(b,1)\} = 010$
 $\{(c,2)\} = 001$
 $\{(b,1), (a,1)\} = 020$
 $\vdots$
 $\{(a,0), (b,1)\} = 110$
 $\vdots$
 $\mathcal{C}_P^\phi = 123$





LET P BE A FINITE POSET AND $\phi: \{0, 1, \dots, m-1\} \rightarrow P$ BE A CHAIN OF P . DENOTE $\mathcal{C}_R^{P, \phi} := \mathcal{I}(\phi(R)) \times \{R\}$, CALLED THE R -COMPONENT, WITH PARTIAL ORDER $(x, R) \leq (y, R)$ IF $x \leq_P y$. DENOTE $\mathcal{C}_P^\phi := \bigsqcup_{0 \leq R < m} \mathcal{C}_R^{P, \phi}$. $\text{TAM}(P, \phi)$ IS THE POSET ORDERED BY INCLUSION ON THE ORDER FILTERS $T \subseteq \mathcal{C}_P^\phi$ SATISFYING FOR ALL $0 \leq i < j < m$, IF $(x, i) \in T$ AND $(\phi(i+1), j) \in T$, THEN $(x, j) \in T$.



ORDER FILTERS:



$\emptyset = 000$
 $\{(a,0)\} = 100$
 $\{(b,1)\} = 010$
 $\{(c,2)\} = 001$
 $\{(b,1), (a,1)\} = 020$
 $\vdots$
 $\{(a,0), (b,1)\} = 110$
 $\vdots$
 $\mathcal{C}_P^\phi = 123$



WHY NOT 110?


SOME RESULTS: DENOTE $L := \text{TAM}(P, \phi)$ AND $C_m := 0 < 1 < \dots < m-1$,

- L IS A LATTICE WITH MEET GIVEN BY INTERSECTION.
- $\ell(L) = |\text{JI}(L)|$, THUS L IS JOIN-EXTREMAL.
- L IS JOIN-SEMI-DISTRIBUTIVE AND WE KNOW WHEN IT IS SEMI-DISTRIBUTIVE.
- L IS JOIN-CONGRUENCE UNIFORM AND WE UNDERSTAND ITS CONGRUENCES.
- L IS LEFT MODULAR, THUS EL-SHELLABLE.
FOR ALL $x \leq y$, $\mu(x, y) \in \{-1, 0, 1\}$.
- $\text{TAM}(C_m, C_m)$ IS THE TAMARI LATTICE OF ORDER $m+1$.




III) HIGHER TORSION CLASSES :

TYPE A AUSLANDER AND NAKAYAMA ALGEBRAS



OSAMU IYAMA


DENOTE $\mathfrak{S}_m^d$ THE POSET OF NON-DECREASING WORDS OF LENGTH d ON $\{0, 1, \dots, m-1\}$ WITH PRODUCT ORDER.



DENOTE $\mathcal{OS}_m^d$ THE POSET OF NON-DECREASING WORDS OF LENGTH d ON $\{0, 1, \dots, m-1\}$ WITH PRODUCT ORDER.

A **TORSION CLASS** IN $\text{mod } A$ IS A SUBCATEGORY CLOSED BY QUOTIENTS AND EXTENSIONS.

Q: $0 \rightarrow 1 \rightarrow \dots \rightarrow m-1$ $A = kQ$ $\text{TORS}(\text{mod } A)$ IS THE TAMARI LATTICE OF ORDER $m+1$.




DENOTE $\mathcal{OS}_m^d$ THE POSET OF NON-DECREASING WORDS OF LENGTH d ON $\{0, 1, \dots, m-1\}$ WITH PRODUCT ORDER.

A **TORSION CLASS** IN $\text{mod } A$ IS A SUBCATEGORY CLOSED BY QUOTIENTS AND EXTENSIONS.

Q: $0 \rightarrow 1 \rightarrow \dots \rightarrow m-1$ $A = kQ$ $\text{TORS}(\text{mod } A)$ IS THE TAMARI LATTICE OF ORDER $m+1$.


A GENERALIZATION IN HIGHER HOMOLOGICAL ALGEBRA WAS DEFINED [JØRGENSEN (16')].



DENOTE $\mathcal{OS}_m^d$ THE POSET OF NON-DECREASING WORDS OF LENGTH d ON $\{0, 1, \dots, m-1\}$ WITH PRODUCT ORDER.

A **TORSION CLASS** IN $\text{mod } A$ IS A SUBCATEGORY CLOSED BY QUOTIENTS AND EXTENSIONS.

Q: $0 \rightarrow 1 \rightarrow \dots \rightarrow m-1$ $A = kQ$ $\text{TORS}(\text{mod } A)$ IS THE TAMARI LATTICE OF ORDER $m+1$.



A GENERALIZATION IN HIGHER HOMOLOGICAL ALGEBRA WAS DEFINED [JØRGENSEN (16')].


THM [A.H.J. K.P.T. (23')]:

LET $\mathcal{M}$ BE A d -CLUSTER TILTING SUBCATEGORY OF $\text{mod } A$. A SUBCATEGORY OF $\mathcal{M}$ IS A d -TORSION CLASS IFF IT IS CLOSED UNDER d -QUOTIENTS AND d -EXTENSIONS. THEY FORM UNDER INCLUSION A COMPLETE LATTICE $d\text{-TORS}(\mathcal{M})$.

DENOTE $\mathcal{OS}_m^d$ THE POSET OF NON-DECREASING WORDS OF LENGTH d ON $\{0, 1, \dots, m-1\}$ WITH PRODUCT ORDER.

A **TORSION CLASS** IN $\text{mod } A$ IS A SUBCATEGORY CLOSED BY QUOTIENTS AND EXTENSIONS.

Q: $0 \rightarrow 1 \rightarrow \dots \rightarrow m-1$ $A = kQ$ $\text{TORS}(\text{mod } A)$ IS THE TAMARI LATTICE OF ORDER $m+1$.



A GENERALIZATION IN HIGHER HOMOLOGICAL ALGEBRA WAS DEFINED [JØRGENSEN (16')].

THM [A.H.J. K.P.T. (23')]:

LET $\mathcal{M}$ BE A d -CLUSTER TILTING SUBCATEGORY OF $\text{mod } A$. A SUBCATEGORY OF $\mathcal{M}$ IS A d -TORSION CLASS IFF IT IS CLOSED UNDER d -QUOTIENTS AND d -EXTENSIONS. THEY FORM UNDER INCLUSION A COMPLETE LATTICE $d\text{-TORS}(\mathcal{M})$.

TWO MAIN EXAMPLES: THE **HIGHER AUSLANDER AND NAKAYAMA ALGEBRAS OF TYPE A**.


BOTH ADMIT RESP. d -CLUSTER TILTING SUBCATEGORIES $\mathcal{M}_m^d$ AND $\mathcal{M}_\ell^d$.

[IYAMA (11')] [O.T. (12)] [J.K. (19')]

DENOTE $\mathcal{O}_m^d$ THE POSET OF NON-DECREASING WORDS OF LENGTH d ON $\{0, 1, \dots, m-1\}$ WITH PRODUCT ORDER.

A **TORSION CLASS** IN $\text{mod } A$ IS A SUBCATEGORY CLOSED BY QUOTIENTS AND EXTENSIONS.

Q: $0 \rightarrow 1 \rightarrow \dots \rightarrow m-1$ $A = kQ$ $\text{TORS}(\text{mod } A)$ IS THE TAMARI LATTICE OF ORDER $m+1$.



A GENERALIZATION IN HIGHER HOMOLOGICAL ALGEBRA WAS DEFINED [JØRGENSEN (16')].

THM [A.H.J.K.P.T. (23')]: LET $\mathcal{M}$ BE A d -CLUSTER TILTING SUBCATEGORY OF $\text{mod } A$. A SUBCATEGORY OF $\mathcal{M}$ IS A d -TORSION CLASS IFF IT IS CLOSED UNDER d -QUOTIENTS AND d -EXTENSIONS. THEY FORM UNDER INCLUSION A COMPLETE LATTICE $d\text{-TORS}(\mathcal{M})$.


TWO MAIN EXAMPLES: THE **HIGHER AUSLANDER AND NAKAYAMA ALGEBRAS OF TYPE A**.

BOTH ADMIT RESP. d -CLUSTER TILTING SUBCATEGORIES $\mathcal{M}_m^d$ AND $\mathcal{M}_\ell^d$. [IYAMA (11')] [J.K. (19')]

ON $\mathcal{O}_m^d$ WE DEFINE $x \rightsquigarrow y$ IF $x_1 \leq y_1 \leq x_2 \leq y_2 \leq \dots \leq x_d \leq y_d$. ON THE WORDS ON $\mathbb{Z}^d$ WE DEFINE $\tau_d(x_1 \dots x_d) = (x_1 - 1) \dots (x_d - 1)$.

THM [A.H.J.K.P.T. (23')]: $T \subseteq \mathcal{O}_m^{d+1}$ IS A d -TORSION CLASS OF $\mathcal{M}_m^d$ IFF FOR ALL $x, y \in \mathcal{O}_m^{d+1}$ WE HAVE BOTH:

- (1) IF $x \leq y$ AND $x_{d+1} = y_{d+1}$, THEN $x \in T$ IMPLIES $y \in T$.
- (2) IF $x \rightsquigarrow \tau_d(y)$ AND $x, y \in T$, THEN $x_1 x_2 \dots x_d y_{d+1} \in T$.

DENOTE $\mathcal{O}_m^d$ THE POSET OF NON-DECREASING WORDS OF LENGTH d ON $\{0, 1, \dots, m-1\}$ WITH PRODUCT ORDER. 

A **TORSION CLASS** IN $\text{mod } A$ IS A SUBCATEGORY CLOSED BY QUOTIENTS AND EXTENSIONS.

Q: $0 \rightarrow 1 \rightarrow \dots \rightarrow m-1$ $A = kQ$ $\text{TORS}(\text{mod } A)$ IS THE TAMARI LATTICE OF ORDER $m+1$.

A GENERALIZATION IN HIGHER HOMOLOGICAL ALGEBRA WAS DEFINED [JØRGENSEN (16')].

THM [A.H.J.K.P.T. (23')]: LET $\mathcal{M}$ BE A d -CLUSTER TILTING SUBCATEGORY OF $\text{mod } A$. A SUBCATEGORY OF $\mathcal{M}$ IS A d -TORSION CLASS IFF IT IS CLOSED UNDER d -QUOTIENTS AND d -EXTENSIONS. THEY FORM UNDER INCLUSION A COMPLETE LATTICE $d\text{-TORS}(\mathcal{M})$.

TWO MAIN EXAMPLES: THE **HIGHER AUSLANDER AND NAKAYAMA ALGEBRAS OF TYPE A**.

BOTH ADMIT RESP. d -CLUSTER TILTING SUBCATEGORIES $\mathcal{M}_m^d$ AND $\mathcal{M}_\ell^d$. [IYAMA (11')] [J.K. (19')]
O.T. (12)

ON $\mathcal{O}_m^d$ WE DEFINE $x \rightsquigarrow y$ IF $x_1 \leq y_1 \leq x_2 \leq y_2 \leq \dots \leq x_d \leq y_d$. ON THE WORDS ON $\mathbb{Z}^d$ WE DEFINE $\tau_d(x_1 \dots x_d) = (x_{d-1}) \dots (x_1)$.

THM [A.H.J.K.P.T. (23')]: $T \subseteq \mathcal{O}_m^{d+1}$ IS A d -TORSION CLASS OF $\mathcal{M}_m^d$ IFF FOR ALL $x, y \in \mathcal{O}_m^{d+1}$ WE HAVE BOTH:

ORDER FILTER $\rightarrow$
IN THE COMPONENTS

- (1) IF $x \leq y$ AND $x_{d+1} = y_{d+1}$, THEN $x \in T$ IMPLIES $y \in T$.
- (2) IF $x \rightsquigarrow \tau_d(y)$ AND $x, y \in T$, THEN $x_1 x_2 \dots x_d y_{d+1} \in T$.


LET $\phi: i \mapsto ii \dots i$.

PROP [S.]: $[(1)+(2)] \Leftrightarrow [(1) + \text{FOR ALL } 0 \leq i < j < m, \text{ IF } (x, i) \in T \text{ AND } (\phi(i+1), j) \in T, \text{ THEN } (x, j) \in T]$.

Denote $\mathcal{O}_m^d$ the poset of non-decreasing words of length d on $\{0, 1, \dots, m-1\}$ with product order. (22)

A torsion class in $\text{mod } A$ is a subcategory closed by quotients and extensions.

Q: $0 \rightarrow 1 \rightarrow \dots \rightarrow m-1$ $A = kQ$ $\text{Tors}(\text{mod } A)$ is the Tamari lattice of order $m+1$.



A generalization in higher homological algebra was defined [Jørgensen (16')].

THM [A.H.J.K.P.T. (23')]:

Let $\mathcal{M}$ be a d -cluster tilting subcategory of $\text{mod } A$. A subcategory of $\mathcal{M}$ is a d -torsion class iff it is closed under d -quotients and d -extensions. They form under inclusion a complete lattice $d\text{-Tors}(\mathcal{M})$.

Two main examples: the higher Auslander and Nakayama algebras of type A.

Both admit resp. d -cluster tilting subcategories $\mathcal{M}_m^d$ and $\mathcal{M}_\ell^d$. [Iyama (11')] [J.K. (19')] O.T. (12)

On $\mathcal{O}_m^d$ we define $x \rightsquigarrow y$ if $x_1 \leq y_1 \leq x_2 \leq y_2 \leq \dots \leq x_d \leq y_d$. On the words on $\mathbb{Z}^d$ we define $\tau_d(x_1 \dots x_d) = (x_{i-1}) \dots (x_{d-1})$.

THM [A.H.J.K.P.T. (23')]:

$T \subseteq \mathcal{O}_m^{d+1}$ is a d -torsion class of $\mathcal{M}_m^d$ iff for all $x, y \in \mathcal{O}_m^{d+1}$ we have both:

ORDER FILTER
IN THE COMPONENTS

- (1) If $x \leq y$ and $x_{d+1} = y_{d+1}$, then $x \in T$ implies $y \in T$.
- (2) If $x \rightsquigarrow \tau_d(y)$ and $x, y \in T$, then $x_1 x_2 \dots x_d y_{d+1} \in T$.

Let $\phi: i \mapsto ii \dots i$.

PROP [S.]: $[(1)+(2)] \Leftrightarrow [(1) + \text{for all } 0 \leq i < j < m, \text{ if } (x, i) \in T \text{ and } (\phi(i+1), j) \in T, \text{ then } (x, j) \in T]$.

THM [S.]: We have $d\text{-Tors}(\mathcal{M}_m^d) \cong \text{Tam}(\mathcal{O}_m^d, \phi)$ and $d\text{-Tors}(\mathcal{M}_\ell^d)$ is a lattice quotient of $d\text{-Tors}(\mathcal{M}_m^d)$.

I) THM.1: A CONGRUENCE UNIFORM LATTICE IS SHELLABLE IFF IT IS EXTREMAL.

II) THM.2: $\text{TAM}(P, \phi)$ IS JOIN- $\left\{ \begin{array}{l} \text{EXTREMAL} \\ \text{SEMIDISTRIBUTIVE} \\ \text{CONGRUENCE UNIFORM} \end{array} \right.$, LEFT MODULAR...

III) USING A COMBINATORIAL DESCRIPTION [A.H.J.K.P.T.(23')] OF THE d -TORSIONS OF $\mathcal{M}_n^d$:

THM.3: $d\text{-TORS}(\mathcal{M}_n^d) \cong \text{TAM}(\mathcal{O}_n^d, \phi)$ AND $d\text{-TORS}(\mathcal{M}_{\underline{d}}^d)$ IS A LATTICE QUOTIENT OF $d\text{-TORS}(\mathcal{M}_n^d)$.

QUESTIONS:

- $|\text{TAM}(\mathcal{O}_n^d, \phi: i \mapsto ii \dots i)| = \bigcirc_{d \rightarrow +\infty} (|\mathcal{J}(\mathcal{O}_n^{d+1})|)$?
- ENUMERATION IN GENERAL OF $\text{TAM}(P, \phi)$?
- DO THE d -TORSION CLASSES IN GENERAL FORM A JOIN-SEMIDISTRIBUTIVE LATTICE ?

TAKE YOUR FAVORITE POSET AND CHAIN IN IT AND TRY IT !

I) THM.1: A CONGRUENCE UNIFORM LATTICE IS SHELLABLE IFF IT IS EXTREMAL.

II) THM.2: $TAM(P, \phi)$ IS JOIN- $\left\{ \begin{array}{l} \text{EXTREMAL} \\ \text{SEMIDISTRIBUTIVE} \\ \text{CONGRUENCE UNIFORM} \end{array} \right.$, LEFT MODULAR...

III) USING A COMBINATORIAL DESCRIPTION [A.H.J.K.P.T.(23')] OF THE d -TORSIONS OF $\mathcal{M}_n^d$:

THM.3: $d\text{-TORS}(\mathcal{M}_n^d) \cong TAM(\mathcal{O}_n^d, \phi)$ AND $d\text{-TORS}(\mathcal{M}_{\frac{n}{2}}^d)$ IS A LATTICE QUOTIENT OF $d\text{-TORS}(\mathcal{M}_n^d)$.

QUESTIONS:

- $|TAM(\mathcal{O}_n^d, \phi: i \mapsto ii \dots i)| = \bigcirc_{d \rightarrow +\infty} (|J(\mathcal{O}_n^{d+1})|)$?
- ENUMERATION IN GENERAL OF $TAM(P, \phi)$?
- DO THE d -TORSION CLASSES IN GENERAL FORM A JOIN-SEMIDISTRIBUTIVE LATTICE ?

TAKE YOUR FAVORITE POSET AND CHAIN IN IT AND TRY IT !

MERCI 😊