List of Publications (1994–2005)

[1] <u>Hideo Kubo</u>, Asymptotic behaviour of solutions to semilinear wave equations with initial data of slow decay, Math. Methods in the Appl. Sci., 17 (1994), 953–970.

[2] <u>Hideo Kubo</u>, Blow-up for semilinear wave equations with initial data of slow decay in low space dimensions, Differential and Integral Equations, 7 (1994), 315–321.

[3] <u>Hideo Kubo</u> and Kôji Kubota, Asymptotic behaviors of radial solutions to semilinear wave equations in odd space dimensions, Hokkaido Math. J., 24 (1995), 9–51.

[4] <u>Hideo Kubo</u> and Kôji Kubota, Asymptotic behaviors of radially symmetric solutions of $\Box u = |u|^p$ for super critical values p in odd space dimensions, Hokkaido Math. J., 24 (1995), 287–336.

[5] <u>Hideo Kubo</u>, On the critical decay and power for semilinear wave equations in odd space dimensions, Discrete and Continuous Dynamical Systems, 2 (1996), 173–190.

[6] <u>Hideo Kubo</u>, Slowly decaying solutions for semilinear wave equations in odd space dimensions, Nonlinear Anal. TMA., 28 (1997), 327–357.

[7] <u>Hideo Kubo</u> and Kôji Kubota, Asymptotic behaviors of radially symmetric solutions of $\Box u = |u|^p$ for super critical values p in even space dimensions, Japanese J. Math., 24 (1998), 191–256.

[8] <u>Hideo Kubo</u> and Masahito Ohta, Critical blowup for systems of semilinear wave equations in low space dimensions, J. Math. Anal. Appl., 240 (1999), 340–360.

[9] <u>Hideo Kubo</u> and Masahito Ohta, Small data blowup for systems of semilinear wave equations with different propagation speeds in three space dimensions, J. Diff. Eqs., 163 (2000), 475–492.

[10] Akira Hoshiga and <u>Hideo Kubo</u>, Global small amplitude solutions of nonlinear hyperbolic systems with a critical exponent under the null condition, SIAM J. Math. Anal., 31 (2000), 486–513.

[11] <u>Hideo Kubo</u> and Kôji Kubota, Asymptotic behavior of classical solutions to a system of semilinear wave equations in low space dimensions, J. Math. Soc. Japan, 53 (2001), 875–912.

[12] Piero D'Ancona, Vladimir Georgiev and <u>Hideo Kubo</u>, Weighted decay estimates for the wave equation, J. Diff. Eqs., 177 (2001), 146–208.

[13] Vladimir Georgiev, Charlotte Heiming and <u>Hideo Kubo</u>, Supercritical semilinear wave equation with non-negative potential, Commun. PDE., 26 (2001), 2267–2303.

[14] <u>Hideo Kubo</u> and Kôji Kubota, Scattering for systems of semilinear wave equations with different speeds of propagation, Advances in Diff. Eqs., 7 (2002), 441–468.

[15] <u>Hideo Kubo</u> and Kotaro Tsugawa, Global solutions and self-similar solutions of the coupled system of semilinear wave equations in three space dimensions, Discrete and Continuous Dynamical Systems, 9 (2003), 471–482.

[16] Akira Hoshiga and <u>Hideo Kubo</u>, Global solvability for systems of nonlinear wave

equations with multiple speeds in two space dimensions, Differential and Integral Equations, 17 (2004), 593–622.

[17] <u>Hideo Kubo</u>, On point-wise decay estimates for the wave equation and their applications, *in* "Dispersive Nonlinear Problems in Mathematical Physics", 123–148, Quaderni di Matematica, 15, Seconda Universitá di Napoli, 2004.

[18] <u>Hideo Kubo</u> and Masahito Ohta, On systems of semilinear wave equations with unequal propagation speeds in three space dimensions, Funkcial. Ekvac., 48 (2005), 65–98.

[19] <u>Hideo Kubo</u> and Masahito Ohta, On the global behavior of classical solutions to coupled systems of semilinear wave equations, *in* "New trends in the theory of hyperbolic equations", 113–211, Operator Theory Adv. and Appl., 159, Birkhäuser Verlag, 2005.

Pubilcations in Proceeding

[1] <u>Hideo Kubo</u> and Kôji Kubota, Asymptotic behaviors of radially symmetric solutions of $\Box u = |u|^p$ for super critical values p in high dimensions, NONLINEAR WAVES, Proceedings of the Fourth MSJ International Research Institute, Vol. I (1996), 239–244.

[2] <u>Hideo Kubo</u> and Kôji Kubota, Asymptotic behaviors of radially symmetric solutions of $\Box u = |u|^p$ for super critical values p in high dimensions, 数理解析研究所講究録 966 "非線形発展方程式とその応用" (1996), 88–94.

[3] <u>Hideo Kubo</u>, Cauchy problem of nonlinear wave equations with small and smooth initial data, 数理解析研究所講究録 1102 "調和解析学と非線形偏微分方程式" (1999), 91–111.

[4] <u>Hideo Kubo</u> and Masahito Ohta, Global existence and blow-up of the classical solutions to systems of semilinear wave equations in three space dimensions, Rend. Istit. Mat. Univ. Trieste, 31 suppl. 2 (2000), 145–168.

[5] Vladimir Georgiev, Charlotte Kerller and <u>Hideo Kubo</u>, Critical exponent for wave equation with potential, Rend. Istit. Mat. Univ. Trieste, 31 suppl. 2 (2000), 103–127.

[6] Piero D'Ancona, Vladimir Georgiev and <u>Hideo Kubo</u>, Weighted Strichartz estimate for the wave equation and low regularity solutions, Rend. Istit. Mat. Univ. Trieste, 31 suppl. 2 (2000), 51–61.

[7] <u>Hideo Kubo</u>, Time-local well-posedness of (1 + 2)-dimensional wave-map type equations and the null condition (Japanese), *in* "On well-posedness and regularity of solutions to partial differential equations (Kyoto, 2001)", Sūrikaisekikenkyūsho Kōkyūroku No. 1284 (2002), 16–31.

[8] <u>Hideo Kubo</u>, On the small data global existence and scattering for systems of semilinear wave equations, in *Hyperbolic problems and related topics*, 219–234, Grad. Ser. Anal., Int. Press, Somerville, MA, 2003.

[9] <u>Hideo Kubo</u>, Coupled system of semilinear wave equations, in *Lecture Notes of Seminario Interdisciplinare di Matematica*, 75–85, Universitá della Basilicata, Dipartmento di Matematica, 2003.

[**10**] <u>Hideo Kubo</u>, 半線型波動方程式系に対する存在定理, 数理解析研究所講究録 1331 短期共同研究 "非線型双曲型方程式系の解の挙動に関する研究" (2003), 50-66. [11] <u>Hideo Kubo</u> and Akira Hoshiga, 準線型波動方程式系に対する存在定理, 数理解析研究 所講究録 1355 短期共同研究"非線形波動および分散型方程式に関する研究"(2004), 1–23.

[12] <u>Hideo Kubo</u>, 波動方程式の外部問題の解の減衰評価について, 数理解析研究所講究 録 1558 短期共同研究"偏微分方程式に対する境界値問題"(2007), 82–105.